

# On solutions of loop quantum cosmology and their thermodynamics properties

H. M. Sadjadi \*

Department of Physics, University of Tehran,  
P. O. B. 14395-547, Tehran 14399-55961, Iran

March 3, 2013

## Abstract

Loop quantum cosmology is considered in inflationary era. A slow rolling scalar field solution with power law potential is presented in the neighborhood of transition time, i.e. when the universe enters inflationary phase from super-inflation era. The second and the generalized second laws of thermodynamics and their validities and violations are discussed and elucidated through some examples.

## 1 Introduction

Application of loop quantum gravity (LQG) to homogenous and isotropic space time, reduces its symmetries and gives rise to loop quantum cosmology (LQC) [1]. In this theory, due to the effective quantum gravitational effects, the big bang singularity, predicted by general relativity, is replaced by a big bounce, after which a super-inflation phase occurs. Afterwards the universe enters a normal inflationary mode.

By super-inflation we mean a period of super-acceleration of the universe during which the time derivative of the Hubble parameter,  $H$ , is positive:  $\dot{H} > 0$ , in contrast to the standard slow roll inflation where  $\dot{H} < 0$ . This super-acceleration can be viewed as a special feature of modified Friedmann equations in effective theory of loop quantum cosmology. Such a phase can also be occurred in general theory of gravity but for example by introducing exotic matter such phantom which violates the null energy condition, or by adding non-minimal coupling terms to the matter action. By considering the super-inflation, the horizon problem may be solved with only a few e-folds [2]. Perturbations in this period of evolution of the universe has also been studied in some papers [3].

---

\*M.sad@ut.ac.ir

As in LQC, the energy density and the Hubble parameter are bounded, the future singularity arisen in dark energy models may also be avoided in this model.

The most adopted source for the matter in inflation scenario is a scalar field with a suitable potential [4]. However, obtaining an exact solution to the modified Friedmann equations in LQC is not possible for a general potential. Inflation, in the context of general relativity, may be driven by a slow rolling scalar field whose parameters are so fine tuned to provide enough number of e-folding according to astrophysical data [4]. The slow roll solution exists also in LQC, and by using numerical methods it is shown the problem of fine tuning is alleviated in this model [5].

To investigate the relationship between thermodynamics and gravity, many studies about thermodynamics properties of cosmological event horizons and thermal nature of the enclosed matter have been done in recent years [6]. In this regard, validity of the second and generalized second laws (GSL) of thermodynamics, in different models of gravity has been subject of many studies [7]. Some of these works have been inspired by the attempts to know the nature of matter with negative pressure such as dark energy and inflaton causing the acceleration or super-acceleration of the universe [8].

The scheme of the paper is as follows. We consider a spatially flat homogeneous and isotropic Friedmann-Lemaître-Robertson-Walker (FLRW) space-time and will focus on inflationary era. After a brief introduction to loop quantum cosmology and one of its exact solutions, i.e. single fluid solution with constant equation of state parameter, we present a slow rolling scalar field solution with power law potential, in the era of transition from super-inflation to inflation phase. We show that this slow rolling solution, in principle exists, provided that some conditions be satisfied. Afterward we study the behavior of the entropy and investigate the validity of the second and generalized second laws of thermodynamics in this model. In our study the pre-bounce era is not considered. We use natural units  $\hbar = c = G = 1$  throughout the paper.

## 2 loop quantum cosmology, preliminaries, solutions

We consider a spatially flat FLRW space-time

$$ds^2 = -dt^2 + a^2(t)(dx^2 + dy^2 + dz^2), \quad (1)$$

where  $a(t)$  is the scale factor. By coupling the matter to classical phase space of FLRW universe, and by taking into account the constraints coming

from loop quantum gravity, one obtains [9]:

$$\begin{aligned}\dot{v} &= \frac{3v}{\lambda\gamma} \sin(\lambda b) \cos(\lambda b) \\ \mathcal{H}_{matt} &= \frac{3}{4\gamma\lambda^2} v \sin^2(\lambda b).\end{aligned}\tag{2}$$

$\mathcal{H}_{matt}$  is the matter Hamiltonian.  $\gamma \approx 0.2375$  denotes the Barbero-Immirzi parameter and  $\lambda$  is the length gap  $\lambda \approx 2.273$ .  $v$  is proportional to the physical volume of a cubical cell, with unit comoving volume :  $v = \frac{a^3}{2\pi\gamma}$ , and  $b$  is the conjugate momentum. Dot denotes derivative with respect to the time "t". The above equations can be rewritten as

$$\begin{aligned}H &= \frac{\sin(2\lambda b)}{2\gamma\lambda} \\ \frac{\sin^2(\lambda b)}{\gamma^2\lambda^2} &= \frac{8\pi}{3}\rho,\end{aligned}\tag{3}$$

resulting the modified Friedmann equation

$$H^2 = \frac{8\pi}{3}\rho\left(1 - \frac{\rho}{\rho_c}\right).\tag{4}$$

$H$  is the Hubble parameter and the matter density,  $\rho$ , has an upper limit  $\rho \leq \rho_c$  where  $\rho_c = \frac{3}{8\pi\gamma^2\lambda^2} \approx 0.409$ . The continuity equation for the matter is

$$\dot{\rho} + 3H(P + \rho) = 0,\tag{5}$$

which by combining with (4), gives the other Friedmann equation

$$\dot{H} = -4\pi(P + \rho)\left(1 - \frac{2\rho}{\rho_c}\right),\tag{6}$$

where  $P$  is the matter pressure. Although for  $\rho \ll \rho_c$  we obtain the usual Friedmann equations, but in the Planck regime, where  $\rho \sim \rho_c$ , the behavior of the system changes drastically. At  $H = 0$ , where  $\rho = \rho_c$ , instead of singularity we have a bounce. If the matter comprising the system satisfies  $P + \rho > 0$ , we have  $\dot{\rho} < 0$  and  $\rho$  decreases after the bounce. For  $\rho_c < \rho < \frac{\rho_c}{2}$ ,  $\dot{H} > 0$  and the universe has a super-inflationary evolution. For  $\rho < \frac{\rho_c}{2}$ , the universe has a normal inflationary mode. Note that if  $\lim_{t \rightarrow \infty} \rho = \frac{\rho_c}{2}$ , the super-inflation lasts forever. As we will show, there may also be a phase of slow roll inflation during both inflation and super-inflation phases. To get some intuition about these behaviors, consider a universe filled with a perfect fluid whose the equation of state parameter,  $\omega$  defined by  $P = \omega\rho$ , is a constant.  $\rho$  is obtained as

$$\rho = \frac{\rho_c}{1 + 6\pi\rho_c(1 + \omega)^2 t^2}.\tag{7}$$

The transition time,  $t_0$  (defined as the time when the super-inflation ends and normal inflation begins), is given by

$$t = \sqrt{\frac{1}{6\pi\rho_c(\omega+1)^2}}. \quad (8)$$

For a massless scalar field,  $\omega = 1$ , and for a fast rolling scalar field  $\omega \simeq 1$ . For radiation  $\omega = \frac{1}{3}$  and for dust or cold dark matter  $\omega = 0$ . Super-inflation lasts longer for smaller  $\omega$  (see fig. (1)).




Figure 1:  $H^2$ , in terms of time for radiation (line), and cold dark matter (points)

Obtaining an exact solution for (4), and (6) is not generally possible. Indeed, (5) can not be easily integrated to give an analytic expression for  $\rho(t)$  (note that only two of the equations (4), (5), and (6) are independent). In the following we present a scalar field solution with power law potential which crosses  $\dot{H}(t_0) = 0$  line, during its slow roll evolution.

## 2.1 Slow roll scalar field solution crossing $\dot{H} = 0$ line

In this part by presenting a series solution to Friedmann equations for a scalar field near the transition time, we show that crossing  $\dot{H} = 0$  is, in principle, possible during slow roll evolution. For simplicity we take  $t_0 = 0$ .

The energy density of the scalar field is

$$\rho = \frac{1}{2}\dot{\phi}^2 + V(\phi) \quad (9)$$

where  $V(\phi)$  is the potential and the pressure is given by

$$P = \frac{1}{2}\dot{\phi}^2 - V(\phi). \quad (10)$$

In the neighborhood of  $t_0 = 0$ , the differentiable Hubble parameter can be presented by [10]

$$H = h_0 + h_1 t^p + \mathcal{O}(t^{p+1}), \quad (11)$$

where  $h_0 = \sqrt{\frac{2\pi\rho_c}{3}}$  is the value of the Hubble parameter at  $t = 0$ , and  $p$  is the order of the first non zero time derivative of  $H$  at  $t = 0$ . If a negative  $h_1 = \frac{1}{p!} \frac{d^p H}{dt^p}(0)$  and an even integer number  $p$  are found such that (11) satisfies the modified Friedmann equations, then it is a solution in the neighborhood of the transition time. Substituting (11) into (12) yields

$$\ddot{\phi} + 3H\dot{\phi} + V'(\phi) = 0, \quad (12)$$

where  $V'(\phi) = \frac{dV(\phi)}{d\phi}$ . By the assumption  $\ddot{\phi} \ll 3H\dot{\phi}$ , we obtain

$$3(h_0 + h_1 t^p + \mathcal{O}(t^{p+1}))\dot{\phi} + V'(\phi) = 0, \quad (13)$$

whose the solution for the power law potential

$$V(\phi) = v_1 \phi^n, \quad (14)$$

is given by

$$\phi = \left( \frac{n(n-2)v_1 t \Phi(-\frac{t^p h_1}{h_0}, 1, \frac{1}{p})}{3h_0 p} + c_1 \right)^{\frac{1}{2-n}}. \quad (15)$$

$c_1$  is determined by

$$c_1 = \phi^{2-n}(t=0), \quad (16)$$

and  $\Phi$  is the Lerchphi function

$$\Phi(x, a, b, ) = \sum_{n=0}^{\infty} \frac{x^n}{(b+n)^a}. \quad (17)$$

For quadratic potential,

$$V(\phi) = \frac{1}{2} m^2 \phi^2, \quad (18)$$

we obtain

$$\phi = \phi(0) \exp \left[ \frac{-m^2 t \Phi(-\frac{t^p h_1}{h_0}, 1, \frac{1}{p})}{3h_0 p} \right]. \quad (19)$$

The slowly varying condition  $\ddot{\phi} \ll 3H\dot{\phi}$ , used to obtain the above solution, is satisfied when

$$\left| \frac{n(n-1)v_1}{9h_0^2} \phi^{n-2}(0) \right| \ll 1, \quad (20)$$

which for quadratic potential gives  $m^2 \ll h_0^2$ , leading to  $m^2 \ll \rho_c$ . Combining (20) with

$$v_1 \phi^n(0) \simeq \frac{\rho_c}{2}, \quad (21)$$

yields

$$\left| \frac{n(n-1)}{12\pi} \right| \ll \phi^2(0). \quad (22)$$

So the value of the scalar field at transition time must be large in this approximation. For  $n \sim \mathcal{O}(1)$ , (22) can be rewritten in terms of  $v_1$  and  $\rho_c$  as  $\left| \left( \frac{\rho_c}{v_1} \right)^{\frac{n}{2}} \right| \ll 1$ .

By inserting (15) into the Friedmann equation

$$\dot{H} = -4\pi\dot{\phi}^2 \left( 1 - \frac{2\rho}{\rho_c} \right), \quad (23)$$

we arrive at

$$\begin{aligned} \dot{H} &= -\frac{8\pi}{27\rho_c} \left( \frac{n^2 v_1^2 \phi^{2(n-1)}(0)}{h_0^3} \right) \left( \frac{n^3(n-1)v_1^3 \phi^{3n-4}(0)}{9h_0^2} + n^2 v_1^2 \phi^{2(n-1)}(0) \right) t \\ &\quad + \mathcal{O}(t^2) \end{aligned} \quad (24)$$

So (11) satisfies Friedmann equation provided that

$$\begin{aligned} h_1 &= -\frac{4\pi}{27\rho_c} \left( \frac{n^2 v_1^2 \phi^{2(n-1)}(0)}{h_0^3} \right) \left( \frac{n^3(n-1)v_1^3 \phi^{3n-4}(0)}{9h_0^2} + n^2 v_1^2 \phi^{2(n-1)}(0) \right), \\ p &= 2. \end{aligned} \quad (25)$$

Applying the slow roll condition (20) to (25), we get

$$h_1 \simeq -\frac{4\pi}{27} \frac{n^4 v_1^4 \phi^{4(n-1)}(0)}{h_0^3 \rho_c}. \quad (26)$$

So this solution describes a transition from super-inflation to inflation at  $t = 0$ , provided that

$$\phi^{4(n-1)}(0) > 0, \quad (27)$$

which is obviously satisfied when  $n$  is an integer. The Hubble parameter and the energy density are then given by

$$\begin{aligned} H &\simeq \sqrt{\frac{2\pi}{3}\rho_c} - \sqrt{\frac{2}{27\pi}} \frac{n^4 v_1^4 \phi^{4(n-1)}(0)}{\rho_c^{\frac{5}{2}}} t^2 + \mathcal{O}(t^3) \\ \rho &\simeq \frac{\rho_c}{2} - \frac{n^2 v_1^2 \phi^{2(n-1)}(0)}{\sqrt{6\pi\rho_c}} t + \mathcal{O}(t^2), \end{aligned} \quad (28)$$

where  $\phi(0)$  is determined by (21). Therefore we conclude that crossing the super-inflation-inflation divide line is, in principle, possible during a slow rolling evolution of a scalar field with power law potential, and the approximate solution to Friedmann equations, in this region, is given by (28).

### 3 Second and generalized second laws

In this part we try to test validity of the generalized second law (GSL) of thermodynamics in a volume enclosed by the apparent horizon in loop quantum cosmology in an expanding FLRW universe. We will elucidate our results with some examples and also the solutions discussed in the previous section.

The entropy of the horizon  $R_h$  in loop quantum gravity is given by [11]

$$S_h = \pi R_h^2 + \pi\alpha \ln(\pi R_h^2) + \beta, \quad (29)$$

where  $\alpha \sim \mathcal{O}(1)$  and  $\beta$  are real constants arisen from quantum corrections.

A natural choice for the cosmological horizon is the apparent horizon  $R_h = \frac{1}{H}$ . Note that  $R_h$  does not exist for  $H = 0$ , but  $R_h(H \neq 0) \in \mathbb{R}$ . By adopting this choice we obtain

$$\dot{S}_h = -2\pi(1 + \alpha H^2) \frac{\dot{H}}{H^3}. \quad (30)$$

Near  $H = 0$ ,  $S_h$  is very large and  $\dot{S}_h \approx -2\pi \frac{\dot{H}}{H^3}$ . So after the bounce,  $S_h$  decreases very fast, reducing  $S_h$  to smaller finite values. In the super accelerated expansion era,  $H$  increases from  $H = 0$  at the bounce to  $H = \sqrt{\frac{2\pi\rho_c}{3}}$  at the transition time, hence  $S_h$  increases only for  $-\frac{3}{2\pi\rho_c} < \alpha < 0$ , and when  $(1 + \alpha H^2) < 0$ . After the transition, we have  $\dot{H} < 0$ , and  $\dot{S}_h > 0$  is valid for  $(1 + \alpha H^2) > 0$ .

To study the GSL, we must also consider the contribution of matter, which satisfies weak energy condition  $\rho > 0$ ,  $P + \rho \geq 0$ , to entropy. We use the first law of thermodynamics

$$dE = TdS_{in} - PdV, \quad (31)$$

to obtain

$$(P + \rho)dV + Vd\rho = TdS_{in}. \quad (32)$$

$T > 0$  is the temperature and  $S_{in}$  is the entropy of the matter inside the horizon. (33) and (5) result in

$$\dot{S}_{in} = -4\pi \frac{(P + \rho)}{TH^2} \left(1 + \frac{\dot{H}}{H^2}\right). \quad (33)$$

During inflation and accelerated expansion,  $1 + \frac{\dot{H}}{H^2} > 0$ , therefore  $\dot{S}_{in} < 0$ . After the inflation  $S_{in}$  decreases.

By adding (33) to (30), we find that GSL is valid whenever

$$\dot{S}_{tot} = -2\pi(1 + \alpha H^2) \frac{\dot{H}}{H^3} - 4\pi \frac{P + \rho}{TH^2} \left(1 + \frac{\dot{H}}{H^2}\right) \geq 0. \quad (34)$$

Provided that the weak energy condition is valid, immediately after the bounce  $\dot{S}_{tot}$  quickly decreases. Afterward, the energy density decreases and the system approaches to the transition time. In the accelerated expansion era ( $\dot{H} + H^2 > 0$ ), validity of GSL requires

$$\dot{H}(1 + \alpha H^2) < 0, \text{ for } \dot{H} > 0, \quad (35)$$

and

$$\dot{H}(1 + \alpha H^2) < 0, \text{ for } \dot{H} < 0. \quad (36)$$

By comparing (35) and (36), and assuming that  $H$  is continuous at transition time, the value of  $\alpha$  is fixed as

$$\alpha = -\frac{1}{H^2(t_0)} = -\frac{3}{2\pi\rho_c} \approx -1.16. \quad (37)$$

But this is not the whole history: In the super accelerated expansion epoch,  $H$  increases until it reaches its maximum value  $H = \sqrt{\frac{2\pi\rho_c}{3}}$  at transition time. But this result is in contradiction with (35) which implies  $H(t < t_0) > \sqrt{\frac{2\pi\rho_c}{3}}$ . So we conclude that, besides a time interval after the bounce, GSL cannot hold in whole of a connected region comprising "a part of super-inflation", transition, and ordinary inflation eras. Now let us have a closer look to the behavior of the system in the neighborhood of transition time  $t_0$ , when  $\dot{S}_{tot}$  reduces to

$$\dot{S}_{tot}(t = t_0) = -4\pi \frac{P + \rho}{TH^2}. \quad (38)$$

The only way to save GSL in this region for ordinary matter (i.e. matter satisfying weak energy condition), is to have  $P(t_0) + \rho(t_0) = 0$ . For the scalar field model,  $P + \rho = \dot{\phi}^2 \geq 0$ . Based on the solutions derived in the previous section, the slow rolling scalar field with power law potential (14) has the following behavior in neighborhood of  $\dot{H} = 0$ :

$$\dot{\phi}^2(t) = \frac{n^2 v_1^2 \phi^{2(n-1)}(t_0)}{6\pi\rho_c} - \frac{2n^3(n-1)v_1^3 \phi^{3n-4}(0)}{\sqrt{216}\pi^{\frac{3}{2}}\rho_c^{\frac{3}{2}}}(t-t_0) + \mathcal{O}((t-t_0)^2). \quad (39)$$

But in the slow roll approximation,  $V(\phi) \approx \frac{\rho_c}{2}$ , then  $\phi(t_0) \neq 0$ , and we have  $\dot{S}_{tot}(t_0) < 0$ . Therefore at  $t_0$  and at least in the neighborhood of  $t_0$  GSL is violated.

At the end of inflation,  $t_{end}$ ,  $\dot{H} + H^2 = 0$ , and

$$\dot{S}_{tot}(t_{end}) = \frac{2\pi}{H}(1 + \alpha H^2). \quad (40)$$

Hence GSL violation is ceased at this time provided that  $1 + \alpha H^2 > 0$  which is always true for  $\alpha > 0$  (a positive value for  $\alpha$  was proposed by [12]).

To be more quantitative and to get more insights about the GSL let us examine (34) using some of the solutions of Friedmann equations. To do this, we have to specify the matter temperature. This temperature is assumed to be equal to the horizon temperature which can be expressed in terms of surface gravity [13],  $\kappa = -H \left(1 + \frac{\dot{H}}{2H^2}\right)$ , as

$$T = \left| \frac{\kappa_H}{2\pi} \right| = \left| \frac{H}{2\pi} \left(1 + \frac{\dot{H}}{2H^2}\right) \right|. \quad (41)$$

So GSL states

$$\dot{S}_{tot} = 2\pi \frac{\dot{H}}{H^3} \left[ -(1 + \alpha H^2) + \left( \frac{H^2 + \dot{H}}{2H^2 + \dot{H}} \right) \left( \frac{2\rho_c}{\rho_c - 2\rho} \right) \right] \geq 0. \quad (42)$$

Hence GSL holds whenever the above inequality is satisfied.

For a barotropic fluid with constant  $w > 0$ , and by considering  $\rho < \rho_c$ , one can show that (42) reduces to

$$(1 - 2x) \left[ -1 - \alpha_c x(1 - x) + \left( \frac{2}{1 - 2x} \right) \left( \frac{-3\Gamma(1 - 2x) + 2(1 - x)}{4(1 - x) - 3\Gamma(1 - 2x)} \right) \right] \leq 0 \quad (43)$$

where  $\Gamma = w + 1$ , and  $x := \frac{\rho}{\rho_c}$ ,  $\alpha_c := \frac{8\pi}{3}\alpha\rho_c$  are dimensionless parameters. According to our previous claim, when  $x \rightarrow 1$ , the GSL is obviously violated. The same occurs when  $x = 1/2$ , independently of the values of the parameters  $\alpha$  and  $\Gamma > 1$ . for  $x \rightarrow 0$  and  $\alpha H^2 \ll 1$ , (42) reduces to  $\pi \frac{\dot{H}^2}{H^3} \geq 0$  which is true. This is the domain of classical theory of gravity far from the Planck era. In this era GSL holds. The inflation ends at  $x = \frac{3\Gamma - 2}{6\Gamma - 2}$ . In this region  $\alpha > 0$  is a sufficient condition for validity of GSL.

In figure (2),  $\dot{S}_{tot}$  is depicted for a barotropic fluid whose equation of state parameter is  $\omega \approx 1$  (e.g. a massless scalar field or a fast rolling scalar field). GSL holds only for  $t > 0.22$  where the inflation nearly ends. The transition time is  $t_0 = 0.18$  (to get more intuition, note these numbers are in terms of the fundamental time unit,  $\sqrt{\frac{\hbar G}{c^5}}$ )

In the above example we assumed that the equation of state parameter is a constant. To get a more complete feel for the behavior of entropy for nonconstant  $w$ , we use numerical method, to depict  $\dot{S}$ , when the inflaton is a scalar field with quadratic potential. We assume that at the bounce the kinetic energy is extremely dominated such that  $\frac{V(\phi)}{\rho_c} \approx 0$ . This can be achieved by suitably choosing the initial values of  $\phi$  and  $\dot{\phi}$  at the bounce. We take  $\phi_B = 1.1$ ,  $\dot{\phi}_B = 0.905$ ,  $m = 6 \times 10^{-7} M_{Pl}$  and  $\alpha = -1.16$ . For  $t < t \sim 0.2138$  GSL fails (see figure (3)).




Figure 2:  $\dot{S}_{tot}$  in terms of time for a single fluid with  $\omega = 1$  and  $\alpha = -1.16$




Figure 3:  $\dot{S}$  in terms of time for a scalar field model with quadratic potential and extreme domination of kinetic energy at the bounce

## 4 Conclusion

We considered loop quantum cosmology, i.e. where the effective quantum corrections coming from loop quantum gravity cannot be neglected. We presented a slow rolling scalar field solution with power law potential in terms of Lerchphi function and showed that a slow rolling scalar field with power law potential can describe the transition from super-inflationary to inflationary phase, provided that the condition (27) be satisfied. The thermodynamics second law and generalized second law (GSL) for the apparent horizon were discussed. If the matter respects the null energy condition, immediately after the bounce the (total) entropy decreases very fast from its huge values, but this decrease does not continue for a long time and after awhile begins to increase. It was shown that the GSL cannot be hold in the whole of a connected region comprising "a part of super-inflation"-transition-inflation epochs. Specially it was shown that for a scalar field this law is violated at transition time. As illustrations, we discussed the behavior of the entropy

for an exactly solvable model (fast roll scalar field) where GSL is violated even after the transition time and holds (nearly) after the inflation ends. Afterwards using numerical method we depicted time derivative of the total entropy for scalar field model whose kinetic energy is extremely dominated at the bounce, and the violation of GSL in a finite time interval after the bounce was demonstrated.

So it seems that GSL does not hold near the bounce where the apparent horizon is very large, and at least, depending on the parameters of the model, this violation may continue until the end of inflation, as was illustrated through some examples. This violation is related to effective quantum effects and GSL continues to be true far from the Planck era. This may be related to the fact that in our computation we have ignored the contribution of the radiation energy density, generated by particle creation from the horizon [14]. Another possibility to alleviate this problem, may be, adoption another horizon such as the future event horizon [15]. Similar violation of GSL was also reported in a super-accelerated epoch in a phantom dominated universe or in modified theories of gravity in the literature [16].

## References

- [1] M. Bojowald, Living Rev. Relativity 8, 11 (2005); A. Ashtekar, M. Bojowald, and J. Lewandowski, Adv. Theor. Math. Phys. 7, 233 (2003); Abhay Ashtekar, Gen. Rel. Grav. 41, 741 (2009); A. Ashtekar, Nuovo Cim. 122B, 135 (2007), arXiv:gr-qc/0702030.
- [2] E. J. Copeland, D. J. Mulryne, N. J. Nunes, and M. Shaeri, Phys. Rev. D 77, 023510 (2008).
- [3] G. M. Hossain, Class. Quant. Grav. 22, 2511 (2005); G. Calcagni and M. Cortes, Class. Quant. Grav. 24, 829 (2007); D. J. Mulryne and N. J. Nunes, Phys. Rev. D 74, 083507 (2006); Y. S. Piao and Y. Z. Zhang, Phys. Rev. D 70, 063513 (2004).
- [4] A. Linde, Particle Physics and Inflationary Cosmology (Harwood, Chur, Switzerland, 1990); A. Linde, Phys. Lett. B 129, 177 (1983).
- [5] A. Ashtekar, D. Sloan, Phys.Lett.B. 694, 108 (2010).
- [6] T. Jacobson, Phys. Rev. Lett. 75, 1260 (1995); R. G. Cai and S. P. Kim, J. High Energy Phys. 02, 050 (2005) 050; R. X. Miao, M. Li, and Y. G. Miao, JCAP 11, 033 (2011); Sh. F. Wu, B. Wang, G. H. Yang, and P. M. Zhang, Class. Quant. Grav. 25, 235018 (2008); K. Bamba, M. Jamil, D. Momeni, and R. Myrzakulov, arXiv:1202.6114v1 [physics.gen-ph]; V. Faraoni, arXiv:1005.2327v1 [gr-qc]; A. Ashtekar and E. Wilson-Ewing, Phys. Rev. D 78, 064047 (2008); K. Karami, A. Abdolmaleki,

- N. Sahraei, and S. Ghaffari, J. High Energy Phys. 1108, 150 (2011); H. M. Sadjadi, Phys. Scripta 05, 055006 (2011); H. M. Sadjadi and M. Jamil, Europhys. Lett 92, 69001 (2010); V. Faraoni, A. F. Z. Moreno, and R. Nandra, arXiv:1202.0719v1 [gr-qc]; K. Karami, M.S. Khaledian, and N. Abdollahi, arXiv:1201.4817v4 [physics.gen-ph]; H. M. Sadjadi, Europhys. Lett. 92, 50014 (2010); U. Debnath, Europhys. Lett. 94, 29001 (2011); V. Faraoni, Phys. Rev. D 80, 044013 (2009); M. Akbar, Int. J. Theor. Phys. 48, 2665 (2009); A. Das, S. Chattopadhyay, and U. Debnath, Found. Phys. 42, 266 (2011); Y. Zhang, Y. Gong, and Z. H. Zhu, Int. J. Mod. Phys. 20, 1505 (2011); H. M. Sadjadi and M. Honardoost, Phys. Lett. B 647, 231 (2007); Z. G. Liu and Y. S. Piao, arXiv:1203.4901 [gr-qc]; Y. S. Piao and E. Zhou, Phys. Rev. D 68, 083515 (2003).
- [7] P. C.W. Davies, Class. Quant. Grav. 4, L255 (1987); 5, 1349 (1988); D. Pavon, Class. Quant. Grav. 7, 487 (1990); R. Brustein, Phys. Rev. Lett. 84, 2072 (2000); T. M. Davis, P. C. W. Davies, and C. H. Lineweaver, Class. Quant. Grav. 20, 2753 (2003); K. Bamba, R. Myrzakulov, S. Nojiri, and S. D. Odintsov, arXiv:1202.4057v3 [physics.gen-ph]].
- [8] G. Izquierdo, D. Pavon, Phys. Lett. B 639, 1 (2006); M. D. Pollock and T. P. Singh, Class. Quant. Grav. 6, 901 (1989); S. Nojiri, S. D. Odintsov, Phys. Rev. D 72, 023003 (2005); H. Farajollahi, A. Salehi, and F. Tayebi, Can. J. Phys. Vol. 89, 915 (2011); S. Nojiri and S. D. Odintsov, Phys. Rev. D 70, 103522 (2004).
- [9] A. Ashtekar, and D. Sloan, Gen. Rel. Grav. 43, 3619 (2011).
- [10] H. M. Sadjadi and M. Alimohammadi, Phys. Rev. D 74, 043506 (2006).
- [11] K. A. Meissner, Class. Quant. Grav. 21, 5245 (2004), A. Ghosh and P. Mitra, Phys. Rev. D 71, 027502 (2005).
- [12] S. Hod, Class. Quant. Grav. 21, L97 (2004).
- [13] R. G. Cai and S. P. Kim, J. High Energy Phys. 0502, 050 (2005).
- [14] S. K. Modak and D. Singleton, arXiv:1205.3404v1 [gr-qc].
- [15] A. Ashtekar and E. Wilson-Ewing, Phys. Rev. D 78, 064047 (2008).
- [16] G. Izquierdo and D. Pavon, Phys. Lett. B 633, 420 (2006); H. Mohseni Sadjadi, Phys. Rev. D 73, 063525 (2006); H. Mohseni Sadjadi, Europhys. Lett. 92, 50014 (2010).